Crupina crupinastrum

Habitus and growth type

Plant height [m]: **0.39**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **12.44**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **9.5**
Moisture indicator value: **2.5**
Reaction indicator value: **7**
Nutrient indicator value: **2.5**
Salinity indicator value: **0.5**

Disturbance indicator values

Disturbance frequency: **0.76**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.82**
Disturbance severity: **0.71**
Disturbance severity (herb layer): **0.52**
Mowing frequency: **0.23**
Grazing pressure: **0.37**
Soil disturbance: **0.25**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FI (TUB) *Helianthemetea guttati***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S63 Eastern garrigue**, **S72 Eastern Mediterranean spiny heath (phrygana)**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz