Cyclamen hederifolium

Habitus and growth type

Plant height [m]: **0.1**
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **July-October**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **3.3**
Temperature indicator value: **8.5**
Moisture indicator value: **4.7**
Reaction indicator value: **6.3**
Nutrient indicator value: **4.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.03**
Disturbance frequency (herb layer): **1.31**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.12**
Mowing frequency: **0**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.22**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CB (PUB) *Quercetea pubescentis***, **FA (QUI) *Quercetea ilicis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T1G Alnus cordata forest**, **T22 Mainland laurophyllous forest**
Constant species of EUNIS habitats: **T1G Alnus cordata forest**, **T22 Mainland laurophyllous forest**

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz