# *Digitalis lanata*

## Habitus and growth type

Plant height [m]: **0.6**
Life span: **Biennial or short-lived, Perennial**
Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.29**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **8.5**
Moisture indicator value: **3**
Reaction indicator value: **8**
Nutrient indicator value: **2**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.32**
Disturbance frequency (herb layer): **1.48**
Disturbance severity: **0.59**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.2**
Grazing pressure: **0.27**
Soil disturbance: **0.14**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CB (PUB) *Quercetea pubescentis***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**

### Broad habitat

Occurrence in broad habitats: **Forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*