Diplotaxis viminea

Habitus and growth type

Plant height [m]: **0.12**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **January-November**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Moisture indicator value: **3.3**
Reaction indicator value: **7.3**
Nutrient indicator value: **6**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **2.04**
Disturbance frequency (herb layer): **2.04**
Disturbance severity: **0.82**
Disturbance severity (herb layer): **0.82**
Mowing frequency: **0.96**
Grazing pressure: **0.22**
Soil disturbance: **0.58**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz