

# ***Dorycnium rectum***

## **Habitus and growth type**

Plant height [m]: **0.9**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

## **Flower**

Flowering period: **March-May**

## **Fruit, seed and dispersal**

Seed mass [mg]: **1.07**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7**

Temperature indicator value: **9**

Moisture indicator value: **7**

Reaction indicator value: **7.3**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.73**

Disturbance frequency (herb layer): **1.54**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.37**

Grazing pressure: **0.22**

Soil disturbance: **0.13**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [CM \(MOL\) Molinio-Arrhenatheretea](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [R31 Mediterranean tall humid inland grassland](#), [T14 Mediterranean and Macaronesian riparian forest](#)

### Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**