Draba nemorosa

Habitus and growth type

Plant height [m]: **0.18**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.04**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Moisture indicator value: **4.3**
Reaction indicator value: **7.3**
Nutrient indicator value: **2.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.79**
Disturbance frequency (herb layer): **1.81**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.53**
Mowing frequency: **0.94**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.32**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CK (SED) *Sedo-Scleranthetea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **8**
Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz