Equisetum palustre

Habitus and growth type

Plant height [m]: **0.37**
Life span: **Perennial**
Life form: **Geophyte**

Leaf

Specific leaf area [mm^2/mg]:

Flower

Flowering period: **March-September**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**
Temperature indicator value: **4.4**
Moisture indicator value: **8.2**
Reaction indicator value: **5.2**
Nutrient indicator value: **3**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.16**
Disturbance frequency (herb layer): **1.71**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.45**
Disturbance severity (herb layer): **0.41**
Mowing frequency: **0.62**
Grazing pressure: **0.19**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***, **IA (ALN) *Alnetea glutinosae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q41 Alkaline, calcareous, carbonate-rich small-sedge spring fen**, **Q42 Extremely rich moss-sedge fen**, **Q43 Tall-sedge base-rich fen**, **Q46 Carpathian travertine fen with halophytes**, **R35 Moist or wet mesotrophic to eutrophic hay meadow**

Constant species of EUNIS habitats: **Q24 Intermediate fen and soft-water spring mire**, **Q41 Alkaline, calcareous, carbonate-rich small-sedge spring fen**, **Q42 Extremely rich moss-sedge fen**, **Q43 Tall-sedge base-rich fen**, **Q45 Arctic-alpine rich fen**, **Q46 Carpathian travertine fen with halophytes**, **Q53 Tall-sedge bed**, **R35 Moist or wet mesotrophic to eutrophic hay meadow**, **R37 Temperate and boreal moist or wet oligotrophic grassland**, **U25 Boreal and Arctic base-rich scree and block field**

Broad habitat

Occurrence in broad habitats: **Aquatic, Spring, Wetland, Mire, Grassland (non-alpine, non-saline), Forest, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **7**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz