Eragrostis cilianensis

Habitus and growth type

Plant height [m]: **0.24**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **24.57**

Flower

Flowering period: **February-October**

Fruit, seed and dispersal

Seed mass [mg]: **0.08**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **8**
Moisture indicator value: **3.2**
Reaction indicator value: **7.3**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **2.22**
Disturbance frequency (herb layer): **2.22**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.85**
Disturbance severity (herb layer): **0.85**
Mowing frequency: **1.13**
Grazing pressure: **0.14**
Soil disturbance: **0.68**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QD (DIG) *Digitario sanguinalis-Eragrostietea minoris***

EUNIS habitat

Diagnostic species of EUNIS habitats: **V34 Trampled xeric grassland with annuals**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz