

# *Erigeron bonariensis*

## Habitus and growth type

Plant height [m]: **0.33**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **13.63**

## Flower

Flowering period: **May-September**

## Fruit, seed and dispersal

Seed mass [mg]: **0.1**

Dispersal mode: **Anemochory, Anthropochory**

Dispersal distance class: **5, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Neophyte**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **7.5**

Temperature indicator value: **9**

Moisture indicator value: **4**

Reaction indicator value: **7**

Nutrient indicator value: **7**

Salinity indicator value: **0.5**

### Disturbance indicator values

Disturbance frequency: **1.24**

Disturbance frequency (herb layer): **1.85**  
Disturbance severity: **0.81**  
Disturbance severity (herb layer): **0.77**  
Mowing frequency: **0.74**  
Grazing pressure: **0.17**  
Soil disturbance: **0.51**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) \*Chenopodietea\*](#), [QD \(DIG\) \*Digitario sanguinalis-Eragrostietea minoris\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T25 \*Phoenix theophrasti\* vegetation](#), [V34 \*Trampled xeric grassland with annuals\*](#)

### Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Sparsely vegetated (incl. rock and scree), Synanthropic**