# *Euphorbia glabriflora*

## Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

## Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **8.2**
Temperature indicator value: **6.7**
Moisture indicator value: **3.7**
Reaction indicator value: **7.1**
Nutrient indicator value: **2.7**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.99**
Disturbance frequency (herb layer): **1.61**
Disturbance severity: **0.38**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.23**
Grazing pressure: **0.29**
Soil disturbance: **0.21**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CK (SED) *Sedo-Scleranthetea*, DA (FES) *Festuco-Brometea***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R17 Heavy-metal dry grassland of the**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

**Balkans, U3A Temperate ultramafic inland cliff**

Constant species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**

Dominant species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*