Fallopia convolvulus

Habitus and growth type

Plant height [m]: **0.48**
Life span: **Annual**
Life form: **Therophyte, Herbaceous liana**

Leaf

Specific leaf area [mm^2/mg]: **22.69**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **4.96**
Dispersal mode: **Myrmecochory, Anthropochory**
Dispersal distance class: **3, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**
Moisture indicator value: **4.9**
Reaction indicator value: **6.2**
Nutrient indicator value: **5.4**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.9**
Disturbance frequency (herb layer): **1.8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.81**
Disturbance severity (herb layer): **0.74**
Mowing frequency: **0.34**
Grazing pressure: **0.14**
Soil disturbance: **0.68**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QA (PAR) *Papaveretea rhoeadis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **V11 Intensive unmixed crops**, **V12 Mixed crops of market gardens and horticulture**, **V13 Arable land with unmixed crops grown by low-intensity agricultural methods**, **V15 Bare tilled, fallow or recently abandoned arable land**
Constant species of EUNIS habitats: **V11 Intensive unmixed crops**, **V12 Mixed crops of market gardens and horticulture**, **V13 Arable land with unmixed crops grown by low-intensity agricultural methods**, **V15 Bare tilled, fallow or recently abandoned arable land**

Broad habitat

Occurrence in broad habitats: **Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz