

# ***Ferulago sylvatica***

## **Habitus and growth type**

Plant height [m]: **0.9**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Flower**

Flowering period: **May-July**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **6.8**

Temperature indicator value: **6.1**

Moisture indicator value: **5.4**

Reaction indicator value: **7**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### **Disturbance indicator values**

Disturbance frequency: **0.52**

Disturbance frequency (herb layer): **1.59**

Disturbance severity: **0.55**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.38**

Grazing pressure: **0.26**

Soil disturbance: **0.14**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) Trifolio-Geranietea sanguinei](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T39 Mediterranean and Balkan subalpine Pinus heldreichii-Pinus peuce forest](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**