Festuca clementei

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Moisture indicator value: **3.6**

Disturbance indicator values

Disturbance frequency: **1.55**
Disturbance frequency (herb layer): **1.55**
Disturbance severity: **0.35**
Disturbance severity (herb layer): **0.35**
Mowing frequency: **0**
Grazing pressure: **0.16**
Soil disturbance: **0.16**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FL (IND) *Festucetea indigestae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1G Iberian oromediterranean siliceous dry grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz