

Galium broterianum

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **7.4**

Temperature indicator value: **6.6**

Moisture indicator value: **8.9**

Reaction indicator value: **5.2**

Nutrient indicator value: **4.8**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **0.67**

Disturbance frequency (herb layer): **1.61**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.03**

Grazing pressure: **0.14**

Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [OD \(PHR\) *Phragmito-Magnocaricetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T22 Mainland laurophyllous forest](#)

Broad habitat

Occurrence in broad habitats: **Wetland**