Helianthemum cinereum

Habitus and growth type

Plant height [m]: **0.15**
Life span: **Perennial**
Life form: **Chamaephyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **5.6**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.54**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **6**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **0.41**
Disturbance frequency (herb layer): **1.49**
Disturbance severity: **0.63**
Disturbance severity (herb layer): **0.24**
Mowing frequency: **0.03**
Grazing pressure: **0.37**
Soil disturbance: **0.15**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **S61 Western basiphilous garrigue**, **S73 Western Mediterranean mountain hedgehog-heath**
Constant species of EUNIS habitats: **S61 Western basiphilous garrigue**, **S73 Western Mediterranean mountain hedgehog-heath**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz