Hippocrepis ciliata

Habitus and growth type

Plant height [m]: **0.09**
Life span: **Annual**
Life form: **Therophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **16.76**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **3.33**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **10**
Moisture indicator value: **2.5**
Reaction indicator value: **8**
Nutrient indicator value: **4.5**
Salinity indicator value: **0.5**

Disturbance indicator values

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency: **1.09**
Disturbance frequency (herb layer): **1.8**
Disturbance severity: **0.76**
Disturbance severity (herb layer): **0.65**
Mowing frequency: **0.1**
Grazing pressure: **0.38**
Soil disturbance: **0.28**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FJ (TRA) *Stipo-Trachynietea distachyae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1F Mediterranean annual-rich dry grassland**
Constant species of EUNIS habitats: **R1F Mediterranean annual-rich dry grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz