# *Hirschfeldia incana*

## Habitus and growth type

Plant height [m]: **0.5**
Life span: **Annual, Biennial or short-lived, Perennial**
Life form: **Hemicryptophyte, Therophyte**

## Leaf

Specific leaf area [$mm^2$/mg]: **19.28**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.25**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.4**
Moisture indicator value: **3**
Reaction indicator value: **7.1**
Nutrient indicator value: **5.6**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.14**
Disturbance frequency (herb layer): **1.95**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance severity: **0.83**
Disturbance severity (herb layer): **0.77**
Mowing frequency: **0.57**
Grazing pressure: **0.21**
Soil disturbance: **0.46**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **QC (CHE) *Chenopodietea***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality amplitude: **5**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*