Hypericum tomentosum

Habitus and growth type

Plant height [m]: **0.48**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **8.7**
Moisture indicator value: **7.5**
Reaction indicator value: **6.5**
Nutrient indicator value: **2.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.17**
Disturbance frequency (herb layer): **2.17**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.49**
Mowing frequency: **0.82**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.34**
Soil disturbance: **0.25**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R32 Mediterranean short moist grassland of lowlands**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Grassland (non-alpine, non-saline), Saline vegetation, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz