Laserpitium halleri

Habitus and growth type

Plant height [m]: **0.45**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **13.32**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **6.83**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **3**
Moisture indicator value: **4.3**
Reaction indicator value: **3**
Nutrient indicator value: **2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.58**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.4**
Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.14**
Grazing pressure: **0.22**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CT (TRI) *Juncetea trifidi***

EUNIS habitat

Diagnostic species of EUNIS habitats: **U34 Mediterranean siliceous inland cliff**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**
Continentality amplitude: **2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz