

Launaea fragilis

Habitus and growth type

Plant height [m]: **0.26**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **11.46**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **12**

Moisture indicator value: **1**

Nutrient indicator value: **1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.51**

Disturbance frequency (herb layer): **1.92**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.37**

Mowing frequency: **0.03**

Grazing pressure: **0.19**

Soil disturbance: **0.51**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) *Ononido-Rosmarinetea*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S65 *Mediterranean gypsum scrub*](#)

Constant species of EUNIS habitats: [S65 *Mediterranean gypsum scrub*](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Scrub**