Lilium martagon

Habitus and growth type

Plant height [m]: **0.78**
Life span: **Perennial**
Life form: **Geophyte**

Leaf

Specific leaf area [mm^2/mg]: **26.09**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Seed mass [mg]: **7.12**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Moisture indicator value: **5.3**
Reaction indicator value: **7.3**
Nutrient indicator value: **5.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.07**
Disturbance frequency (herb layer): **1.27**
Disturbance severity: **0.67**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.14**
Mowing frequency: **0.02**
Grazing pressure: **0.19**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3C Taxus baccata forest**, **T3H Larix light taiga**
Constant species of EUNIS habitats: **T17 Fagus forest on non-acid soils**, **T1F Ravine forest**, **T3C Taxus baccata forest**, **T3H Larix light taiga**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest**

Distribution

Continentality: **6**
Continentality amplitude: **5**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz