Limonium graecum

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **11.5**
Reaction indicator value: **9**
Nutrient indicator value: **6.5**
Salinity indicator value: **5.5**

Disturbance indicator values

Disturbance frequency: **0.76**
Disturbance frequency (herb layer): **0.79**
Disturbance severity: **0.34**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.07**
Grazing pressure: **0.1**
Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JB (CRI) *Crithmo-Staticetea***, **MG (SAL) *Salicornietea fruticosae***

EUNIS habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of EUNIS habitats: **N32 Mediterranean and Black Sea rocky sea cliff and shore**

Broad habitat

Occurrence in broad habitats: **Coastal cliff, Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz