Limonium meyeri

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.53**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **5.3**
Moisture indicator value: **4.5**

Disturbance indicator values

Disturbance frequency: **1.7**
Disturbance frequency (herb layer): **1.91**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.48**
Mowing frequency: **0.09**
Grazing pressure: **0.33**
Soil disturbance: **0.26**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **JB (CRI) *Crithmo-Staticetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **MA241 Black Sea littoral saltmarsh**, **N1E Black Sea broad-leaved coastal dune forest**, **N35 Mediterranean and Black Sea**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

soft sea cliff, **S94 Semi-desert riparian scrub**

Constant species of EUNIS habitats: **MA241 Black Sea littoral saltmarsh**, **N1E Black Sea broad-leaved coastal dune forest**, **N35 Mediterranean and Black Sea soft sea cliff**, **S94 Semi-desert riparian scrub**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal cliff, Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz