Lonicera tatarica

Habitus and growth type

Plant height [m]: **2.13**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Leaf

Leaf type (woody plants only): **Broad deciduous or semi-deciduous**
Leaf phenology (woody plants only): **Winter deciduous**

Fruit, seed and dispersal

Seed mass [mg]: **3.18**
Dispersal mode: **Endozoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **6.5**
Temperature indicator value: **6.8**
Moisture indicator value: **6**

Disturbance indicator values

Disturbance frequency: **0.29**
Disturbance frequency (herb layer): **1.34**
Disturbance severity: **0.67**
Disturbance severity (herb layer): **0.27**
Mowing frequency: **0.1**
Grazing pressure: **0.19**
Soil disturbance: **0.16**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **HA (POP) *Alno glutinosae-Populetea albae***

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz