# *Medicago medicaginoides*

### Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

### Fruit, seed and dispersal

Seed mass [mg]: **0.56**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Disturbance indicator values

Disturbance frequency: **2.18**
Disturbance frequency (herb layer): **2.1**
Disturbance severity: **0.37**
Disturbance severity (herb layer): **0.37**
Mowing frequency: **0.08**
Grazing pressure: **0.33**
Soil disturbance: **0.23**

### Habitat and sociology

#### Syntaxon

Diagnostic species of phytosociological classes: **EA (LER) *Artemisietea lerchianae***

#### EUNIS habitat

Diagnostic species of EUNIS habitats: **S67 Aralo-Caspian semi-desert**
Constant species of EUNIS habitats: **S67 Aralo-Caspian semi-desert**

#### Broad habitat

Occurrence in broad habitats: **Semi-desert**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*