# *Mibora minima*

## Habitus and growth type

Plant height [m]: **0.08**
Life span: **Annual**
Life form: **Therophyte**

## Flower

Flowering period: **February-May**

## Fruit, seed and dispersal

Seed mass [mg]: **0.13**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **1, 7**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **8.3**
Moisture indicator value: **2**
Nutrient indicator value: **2.3**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **2**
Disturbance frequency (herb layer): **2.18**
Disturbance severity: **0.43**
Disturbance severity (herb layer): **0.43**
Mowing frequency: **0.35**
Grazing pressure: **0.29**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Soil disturbance: **0.46**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CJ (COR) *Koelerio-Corynephoretea canescentis*, FI (TUB) *Helianthemetea guttati*, QA (PAR) *Papaveretea rhoeadis***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R1R Mediterranean to Atlantic open, dry, acid and neutral grassland**

### Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

## Distribution

Continentality: **2**
Continentality amplitude: **2**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*