

Muscari neglectum

Habitus and growth type

Plant height [m]: **0.16**

Life span: **Perennial**

Life form: **Geophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **12.54**

Flower

Flowering period: **February-June**

Fruit, seed and dispersal

Seed mass [mg]: **3.17**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **7.1**

Moisture indicator value: **3.4**

Reaction indicator value: **7**

Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.86**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.52**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.27**

Grazing pressure: **0.33**

Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FP \(ONO\) Festuco hystrixis-Ononidetea striatae](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#)

Constant species of EUNIS habitats: [R14 Perennial rocky grassland of the Italian Peninsula](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **8**