# *Odontites luteus*

## Habitus and growth type

Plant height [m]: **0.31**
Life span: **Annual**
Life form: **Therophyte**

## Leaf

Specific leaf area [$mm^2$/mg]: **21.64**

## Flower

Flowering period: **July-October**

## Fruit, seed and dispersal

Seed mass [mg]: **0.22**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **6.8**
Moisture indicator value: **3**
Reaction indicator value: **8.8**
Nutrient indicator value: **1.8**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.54**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Disturbance frequency (herb layer): **1.65**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.34**
Mowing frequency: **0.24**
Grazing pressure: **0.31**
Soil disturbance: **0.16**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

## Distribution

Continentality: **6**
Continentality amplitude: **6**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*