

Ononis reclinata

Habitus and growth type

Plant height [m]: **0.11**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.76**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **9.2**

Moisture indicator value: **1.8**

Reaction indicator value: **7.3**

Salinity indicator value: **0.7**

Disturbance indicator values

Disturbance frequency: **0.87**

Disturbance frequency (herb layer): **1.74**

Disturbance severity: **0.76**

Disturbance severity (herb layer): **0.55**

Mowing frequency: **0.15**

Grazing pressure: **0.35**

Soil disturbance: **0.28**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#),
[S63 Eastern garrigue](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#)

Constant species of EUNIS habitats: [R1F Mediterranean annual-rich dry grassland](#),
[S63 Eastern garrigue](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**