Ononis striata

Habitus and growth type

Plant height [m]: **0.1**
Life span: **Perennial**
Life form: **Chamaephyte**

Leaf

Specific leaf area [mm^2/mg]: **17.02**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **6**
Moisture indicator value: **3**
Reaction indicator value: **8**
Nutrient indicator value: **1.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.03**
Disturbance frequency (herb layer): **1.71**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.44**
Mowing frequency: **0.57**
Grazing pressure: **0.41**
Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FP (ONO) *Festuco hystricis-Ononidetea striatae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe**, **R1H Iberian oromediterranean basiphilous dry grassland**

Constant species of EUNIS habitats: **R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz