Ononis tridentata

Habitus and growth type

Plant height [m]:
Life span: **Perennial**
Life form: **Chamaephyte**
Spinescence: **not spinescent**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **9**
Moisture indicator value: **2.2**
Reaction indicator value: **8**
Nutrient indicator value: **3.3**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.26**
Disturbance frequency (herb layer): **1.76**
Disturbance severity: **0.39**
Disturbance severity (herb layer): **0.25**
Mowing frequency: **0.02**
Grazing pressure: **0.3**
Soil disturbance: **0.19**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FB (ROS) *Ononido-Rosmarinetea***

EUNIS habitat

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Diagnostic species of EUNIS habitats: **S65 Mediterranean gypsum scrub**
Constant species of EUNIS habitats: **S65 Mediterranean gypsum scrub**
Dominant species of EUNIS habitats: **S65 Mediterranean gypsum scrub**

Broad habitat

Occurrence in broad habitats: **Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz