Ononis verae

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **11**
Moisture indicator value: **3**
Reaction indicator value: **8**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.53**
Disturbance frequency (herb layer): **1.62**
Disturbance severity: **0.77**
Disturbance severity (herb layer): **0.46**
Mowing frequency: **0.17**
Grazing pressure: **0.27**
Soil disturbance: **0.25**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FJ (TRA) *Stipo-Trachynietea distachyae***

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz