Paeonia broteroi

Habitus and growth type

Life span: **Perennial**
Life form: **Geophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7**
Moisture indicator value: **1**
Reaction indicator value: **8**
Nutrient indicator value: **1**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.06**
Disturbance frequency (herb layer): **1.33**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.14**
Mowing frequency: **0.02**
Grazing pressure: **0.26**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FA (QUI) *Quercetea ilicis***

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz