Parentucellia viscosa

Habitus and growth type

Plant height [m]: **0.33**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **24.21**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.02**
Dispersal mode: **Anemochory**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8**
Reaction indicator value: **6**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.18**
Disturbance frequency (herb layer): **1.75**
Disturbance severity: **0.68**
Disturbance severity (herb layer): **0.64**
Mowing frequency: **0.44**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.26**
Soil disturbance: **0.34**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FJ (TRA) *Stipo-Trachynietea distachyae*, QC (CHE) *Chenopodietea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz