

Paronychia argentea

Habitus and growth type

Plant height [m]: **0.03**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **15.68**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.74**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **9.3**

Moisture indicator value: **2.7**

Reaction indicator value: **7**

Nutrient indicator value: **3.5**

Salinity indicator value: **1.7**

Disturbance indicator values

Disturbance frequency: **1.22**

Disturbance frequency (herb layer): **1.99**

Disturbance severity: **0.64**

Disturbance severity (herb layer): **0.59**

Mowing frequency: **0.2**

Grazing pressure: **0.32**

Soil disturbance: **0.41**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**FH \(BUL\) *Poetea bulbosae***](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R1D Mediterranean closely grazed dry grassland**](#)

Broad habitat

Occurrence in broad habitats: **Coastal beach, dune or shingle, Grassland (non-alpine, non-saline), Scrub**