Paronychia macedonica

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.8**
Temperature indicator value: **7.1**
Moisture indicator value: **2.8**
Reaction indicator value: **7.3**
Nutrient indicator value: **2.2**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.14**
Disturbance frequency (herb layer): **2.14**
Disturbance severity: **0.32**
Disturbance severity (herb layer): **0.32**
Mowing frequency: **0.01**
Grazing pressure: **0.32**
Soil disturbance: **0.13**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R45 Alpine and subalpine calcareous grassland of the Balkans and Apennines**

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz