# *Phalacrocarpum oppositifolium*

## Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte, Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **1.67**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **4**
Moisture indicator value: **3**
Reaction indicator value: **3**
Nutrient indicator value: **1**

### Disturbance indicator values

Disturbance frequency: **0.78**
Disturbance frequency (herb layer): **1.4**
Disturbance severity: **0.4**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.05**
Grazing pressure: **0.25**
Soil disturbance: **0.11**

## Habitat and sociology

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R1G Iberian oromediterranean siliceous dry grassland**
Constant species of EUNIS habitats: **R1G Iberian oromediterranean siliceous dry grassland**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*