Pinus brutia

Habitus and growth type

Plant height [m]: **17.55**
Life span: **Perennial**
Life form: **Phanerophyte, Tree**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **44.11**
Dispersal mode: **Anemochory**
Dispersal distance class: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.9**
Moisture indicator value: **3**
Reaction indicator value: **7**
Nutrient indicator value: **2.7**
Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **0.13**
Disturbance frequency (herb layer): **1.54**
Disturbance severity: **0.7**
Disturbance severity (herb layer): **0.19**
Mowing frequency: **0.02**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.24**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CB (PUB) *Quercetea pubescentis*, FA (QUI) *Quercetea ilicis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T3A Mediterranean lowland to submontane Pinus forest**
Constant species of EUNIS habitats: **T3A Mediterranean lowland to submontane Pinus forest**
Dominant species of EUNIS habitats: **T3A Mediterranean lowland to submontane Pinus forest**

Broad habitat

Occurrence in broad habitats: **Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz