

Pistacia lentiscus

Habitus and growth type

Plant height [m]: **3.75**

Life span: **Perennial**

Life form: **Phanerophyte, Tree, Shrub**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **5.44**

Flower

Flowering period: **February-May**

Fruit, seed and dispersal

Seed mass [mg]: **27.04**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Temperature indicator value: **9.5**

Moisture indicator value: **3**

Reaction indicator value: **6.3**

Nutrient indicator value: **4.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.23**
 Disturbance frequency (herb layer): **1.37**
 Disturbance severity: **0.7**
 Disturbance severity (herb layer): **0.25**
 Mowing frequency: **0.01**
 Grazing pressure: **0.28**
 Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FA \(QUI\) *Quercetea ilicis*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N1B Mediterranean and Black Sea coastal dune scrub](#), [N1G Mediterranean coniferous coastal dune forest](#), [S51 Mediterranean maquis and arborescent matorral](#), [S54 Thermomediterranean arid scrub](#), [S62 Western acidophilous garrigue](#), [S71 Western Mediterranean spiny heath](#), [T24 Olea europaea-Ceratonía siliqua forest](#), [T25 Phoenix theophrasti vegetation](#), [T3A Mediterranean lowland to submontane Pinus forest](#)

Constant species of EUNIS habitats: [N1B Mediterranean and Black Sea coastal dune scrub](#), [N1G Mediterranean coniferous coastal dune forest](#), [S51 Mediterranean maquis and arborescent matorral](#), [S53 Spartium junceum scrub](#), [S54 Thermomediterranean arid scrub](#), [S62 Western acidophilous garrigue](#), [S63 Eastern garrigue](#), [S71 Western Mediterranean spiny heath](#), [S72 Eastern Mediterranean spiny heath \(phrygana\)](#), [S93 Mediterranean riparian scrub](#), [T1A Mediterranean thermophilous deciduous forest](#), [T21 Mediterranean evergreen Quercus forest](#), [T24 Olea europaea-Ceratonía siliqua forest](#), [T25 Phoenix theophrasti vegetation](#), [T3A Mediterranean lowland to submontane Pinus forest](#)

Dominant species of EUNIS habitats: [N1B Mediterranean and Black Sea coastal dune scrub](#), [N1G Mediterranean coniferous coastal dune forest](#), [S51 Mediterranean maquis and arborescent matorral](#), [T24 Olea europaea-Ceratonía siliqua forest](#), [T3A Mediterranean lowland to submontane Pinus forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest, Sparsely vegetated (incl. rock and scree)**