# *Quercus canariensis*

## Habitus and growth type

Plant height [m]:
Life span: **Perennial**
Life form: **Phanerophyte, Tree**

## Leaf

Specific leaf area [$mm^2$/mg]: **8.6**

## Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Dyszoochory**
Dispersal distance class: **6**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**
Temperature indicator value: **7**
Moisture indicator value: **5**
Reaction indicator value: **5**
Nutrient indicator value: **5**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.12**
Disturbance frequency (herb layer): **1.41**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.23**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0**
Grazing pressure: **0.21**
Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **FA (QUI) *Quercetea ilicis***

### Broad habitat

Occurrence in broad habitats: **Forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*