Satureja coerulea

Habitus and growth type

Life span: **Perennial**
Life form: **Chamaephyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.32**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **6.6**
Moisture indicator value: **3.6**
Reaction indicator value: **7.5**
Nutrient indicator value: **2.8**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **1.41**
Disturbance frequency (herb layer): **2.07**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.51**
Mowing frequency: **0.65**
Grazing pressure: **0.42**
Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz