

Scorzoneroides cantabrica

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **3**

Moisture indicator value: **3**

Reaction indicator value: **2**

Nutrient indicator value: **1**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.34**

Disturbance frequency (herb layer): **1.55**

Disturbance severity: **0.35**

Disturbance severity (herb layer): **0.32**

Mowing frequency: **0.08**

Grazing pressure: **0.27**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**FL \(IND\) Festucetea indigestae**](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland**