Scrophularia umbrosa

Habitus and growth type

Plant height [m]: **0.67**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **31.83**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.06**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.7**
Moisture indicator value: **8.6**
Reaction indicator value: **7.5**
Nutrient indicator value: **7**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.71**
Disturbance frequency (herb layer): **1.46**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity: **0.46**
Disturbance severity (herb layer): **0.36**
Mowing frequency: **0.28**
Grazing pressure: **0.14**
Soil disturbance: **0.12**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Wetland, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **6**
Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz