# *Sedum urvillei*

## Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Dystrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **8**
Moisture indicator value: **1**
Reaction indicator value: **9**
Nutrient indicator value: **1**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.3**
Disturbance frequency (herb layer): **1.95**
Disturbance severity: **0.52**
Disturbance severity (herb layer): **0.49**
Mowing frequency: **0.33**
Grazing pressure: **0.34**
Soil disturbance: **0.26**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CK (SED) *Sedo-Scleranthetea***

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*