Selinum carvifolia

Habitus and growth type

Plant height [m]: **0.57**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **18.7**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.36**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6.9**
Temperature indicator value: **5**
Moisture indicator value: **7.2**
Reaction indicator value: **6**
Nutrient indicator value: **3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.12**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.84**
Disturbance severity: **0.49**
Disturbance severity (herb layer): **0.45**
Mowing frequency: **0.88**
Grazing pressure: **0.2**
Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea*, IA (ALN) *Alnetea glutinosae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R37 Temperate and boreal moist or wet oligotrophic grassland**
Constant species of EUNIS habitats: **R37 Temperate and boreal moist or wet oligotrophic grassland**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Mire, Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **5**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz