Solidago canadensis

Habitus and growth type

Plant height [m]: **1.2**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm^2/mg]: **25.87**

Flower

Flowering period: **June-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.07**
Dispersal mode: **Anemochory, Anthropochory**
Dispersal distance class: **5, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **6**
Moisture indicator value: **6**
Reaction indicator value: **6.4**
Nutrient indicator value: **6.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.86**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.48**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.57**
Mowing frequency: **0.59**
Grazing pressure: **0.17**
Soil disturbance: **0.24**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QF (ART) *Artemisietea vulgaris*, QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz