Stachys scardica

Habitus and growth type

Life span: **Perennial**
Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Ellenberg-type indicator values

Light indicator value: **6.6**
Temperature indicator value: **6.4**
Moisture indicator value: **4.9**
Reaction indicator value: **5.9**
Nutrient indicator value: **3.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.24**
Disturbance frequency (herb layer): **1.39**
Disturbance severity: **0.58**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.14**
Grazing pressure: **0.24**
Soil disturbance: **0.12**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CK (SED) *Sedo-Scleranthetea*, CO (ERI) *Erico-Pinetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Constant species of EUNIS habitats: **R17 Heavy-metal dry grassland of the Balkans**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz