

Styrax officinalis

Habitus and growth type

Plant height [m]: **4.75**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **411.55**

Dispersal mode: **Endozoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **9**

Reaction indicator value: **6.7**

Nutrient indicator value: **3.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.08**

Disturbance frequency (herb layer): **1.37**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.17**

Mowing frequency: **0**

Grazing pressure: **0.25**

Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T38 Mediterranean montane Cedrus forest](#),
[T3A Mediterranean lowland to submontane Pinus forest](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Forest**