Symphytum asperum

Habitus and growth type

Plant height [m]: **0.96**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **4.86**
Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.8**
Moisture indicator value: **5.7**
Reaction indicator value: **7**
Nutrient indicator value: **8.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.11**
Disturbance frequency (herb layer): **1.39**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.22**
Mowing frequency: **0.02**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.2**
Soil disturbance: **0.11**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **QG (EPI) *Epilobietea angustifolii***

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline), Scrub, Synanthropic**

Distribution

Continentality: **7**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz