Tamarix hampeana

Habitus and growth type

Plant height [m]: **6.87**
Life span: **Perennial**
Life form: **Phanerophyte, Shrub**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8**
Temperature indicator value: **11**
Moisture indicator value: **8**
Reaction indicator value: **9**
Nutrient indicator value: **7**
Salinity indicator value: **6**

Disturbance indicator values

Disturbance frequency: **1.28**
Disturbance frequency (herb layer): **1.63**
Disturbance severity: **0.53**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **0**
Grazing pressure: **0.27**
Soil disturbance: **0.2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FD (NER) *Nerio-Tamaricetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **S93 Mediterranean riparian scrub**

Constant species of EUNIS habitats: **S93 Mediterranean riparian scrub**

Dominant species of EUNIS habitats: **S93 Mediterranean riparian scrub**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Saline vegetation, Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz